

Study of scalar meson $a_0(980)$ from $B \rightarrow a_0(980)\pi$ Decays

Zhi-Qing Zhang^{a,b*} and Zhen-jun Xiao^{b †}

*a Department of Physics, Henan University of Technology,
Zhengzhou, Henan 450052, P.R.China;*

*b Department of Physics and Institute of Theoretical Physics,
Nanjing Normal University, Nanjing, Jiangsu 210097, P.R.China*

Abstract

In this paper, we calculate the branching ratios and the direct CP-violating asymmetries for decays $\bar{B}^0 \rightarrow a_0^0(980)\pi^0, a_0^+(980)\pi^-, a_0^-(980)\pi^+$ and $B^- \rightarrow a_0^0(980)\pi^-, a_0^-(980)\pi^0$ by employing the perturbative QCD factorization approach. Although the light scalar meson $a_0(980)$ is widely perceived as primarily the four-quark bound states, the calculation shows that the 2-quark model supposition for $a_0(980)$ can not be ruled out by the currently available experiment upper limits. In these considered decays, the branching ratio for the $B^- \rightarrow a_0^0(980)\pi^-$ is the largest, 2.8×10^{-6} , while its direct CP asymmetry is the smallest, $\sim 14\%$. Although the direct CP asymmetries for the decays $\bar{B}^0 \rightarrow a_0^0(980)\pi^0, B^- \rightarrow a_0^-(980)\pi^0$ are large, about $(70 \sim 80)\%$, it is still difficult to measure them, since their branching ratios are small, around $(4 \sim 5) \times 10^{-7}$.

PACS numbers: 13.25.Hw, 12.38.Bx, 14.40.Nd

* Electronic address: zhangzhiqing@zzu.edu.cn

† Electronic address: xiaozhenjun@njnu.edu.cn

I. INTRODUCTION

The study about scalar meson is an interesting topic for both theory and experiment. In order to uncover their mysterious structure, intensive studies have been done for the B meson decays involving a scalar meson as one of the two final state mesons. Such kind of decays have been studied by employing various factorization approaches, such as the generalized factorization approach [1], the QCD factorization (QCDF) approach [2, 3], the perturbative QCD (pQCD) approach [4, 5, 6, 7, 8], and by using the QCD sum rule [9, 10, 11].

On the experimental side, from the first scalar meson $f_0(980)$ observed in the decay mode $B \rightarrow f_0(980)K$ by Belle [12], and confirmed by BaBar [13] later, then many channels involving a scalar in the final state have been measured by Belle [14, 15] and BaBar [16, 17]. For example, the decays $B \rightarrow a_0(980)\pi$ were searched by BarBar five years ago [18], especially the decay $B^- \rightarrow a_0^-(980)\pi^0$, which has been considered as a best candidate to distinguish the nature of the scalar $a_0(980)$ [19]. The authors of Ref.[19] argued that if the branching ratio of this channel can be measured accurately by the experiment, one can separate the four- and two-quark assignments, because the results of these two assignments have a difference of one order of magnitude. So in the past three years, BarBar have given this channel twice measurements [20, 21] and get two almost identical upper limits. For our considered decays, only the experimental upper limits are available now for some of them [22]:

$$\begin{aligned} Br(\bar{B}^0 \rightarrow a_0^+(980)\pi^-) &< 3.1 \times 10^{-6}, \\ Br(B^- \rightarrow a_0^-(980)\pi^0) &< 1.4 \times 10^{-6}, \\ Br(B^- \rightarrow a_0^0(980)\pi^-) &< 5.8 \times 10^{-6}. \end{aligned} \tag{1}$$

In this paper, we will study the branching ratios and CP asymmetries of $\bar{B}^0 \rightarrow a_0^0(980)\pi^0, a_0^\pm(980)\pi^\mp$ and $B^- \rightarrow a_0^-(980)\pi^0, a_0^0(980)\pi^-$ within perturbative QCD approach based on k_T factorization. In the following, we use a_0 to denote $a_0(980)$ in some places for convenience. It is organized as follows. In Sect.II, the status of the study on the physical properties of a_0 , the relevant decay constants and light-cone distribution amplitudes are discussed. In Sec.III, we then analysis these decay channels using the pQCD approach. The numerical results and the discussions are given in the section IV. The conclusions are presented in the final part.

II. PHYSICAL PROPERTIES OF THE FINAL PARTICLES

Many scalar mesons below 2GeV have been found in experiments. We can't accommodate these scalar mesons into one nonet, but need at least two nonets below and above 1 GeV[23]. Among them, the scalar mesons below 1 GeV, including $f_0(600)(\sigma), f_0(980), K_0^*(800)(\kappa)$ and $a_0(980)$, are usually viewed to form an SU(3) nonet; while scalar mesons above 1 GeV, including $f_0(1370), f_0(1500)/f_0(1700), K^*(1430)$ and $a_0(1450)$ form the other SU(3) nonet. There are several different scenarios to describe these mesons in the quark model [24, 25, 26, 27]. For example $a_0(980)$ meson, which has been suggested as $\bar{q}q$ lowest lying state [24](called scenario I) or four-quark bound state [25](called scenario II). In the scenario I, the former SU(3) nonet mesons are treated as

the $\bar{q}q$ ground states, and the latter nonet ones are the first excited states; in the scenario II, the former nonet mesons are viewed as four-quark bound states, while the latter nonet ones are $\bar{q}q$ ground states. Some people also consider that it is not made of one simple component but might have a more complex nature such as having a $K\bar{K}$ component[26, 27], even the superpositions of the two- and four- quark states. In order to make quantitative predictions, we identify $a_0(980)$ as the two-quark state in the calculation.

In 2-quark model, the decay constants for scalar meson a_0 are defined by:

$$\langle a_0(p) | \bar{q}_2 \gamma_\mu q_1 | 0 \rangle = f_{a_0} p_\mu, \quad \langle a_0(p) | \bar{q}_2 q_1 | 0 \rangle = m_{a_0} \bar{f}_{a_0}. \quad (2)$$

For the neutral scalar meson a_0 cannot be produced via the vector current (restricted by the charge conjugation invariance or the G parity conservation), the vector decay constant $f_{a_0} = 0$. As to the charged scalar mesons a_0^- , from the equation of motion:

$$\mu_{a_0^-} f_{a_0^-} = \bar{f}_{a_0^-}, \quad \text{with} \quad \mu_{a_0^-} = \frac{m_{a_0^-}}{m_d(\mu) - m_u(\mu)}, \quad (3)$$

its vector decay constant is proportional to the mass difference between the constituent u and d quarks. It is easy to see the vector decay constant is very small, and will equal to zero in the SU(3) limit. So we only need consider the scalar decay constant $\bar{f}_{a_0}$, which is scale dependent. Fixing the scale at 1 GeV, the value is $\bar{f}_{a_0} = (365 \pm 20) \text{ MeV}$, which is calculated in QCD sum rules[3].

The light-cone distribution amplitudes (LCDAs) for the scalar meson a_0 can be written as:

$$\begin{aligned} \langle a_0(p) | \bar{q}_1(z)_l q_2(0)_j | 0 \rangle &= \frac{1}{\sqrt{2N_c}} \int_0^1 dx e^{ixp \cdot z} \\ &\times \{ \not{p} \Phi_{a_0}(x) + m_{a_0} \Phi_{a_0}^S(x) + m_{a_0} (\not{n}_+ \not{n}_- - 1) \Phi_{a_0}^T(x) \}_{jl}, \end{aligned} \quad (4)$$

where n_+ and n_- are light-like vectors: $n_+ = (1, 0, 0_T)$, $n_- = (0, 1, 0_T)$, and n_+ is parallel with the moving direction of the scalar meson a_0 . The normalization can be related to the decay constants:

$$\int_0^1 dx \Phi_{a_0}(x) = \int_0^1 dx \Phi_{a_0}^T(x) = 0, \quad \int_0^1 dx \Phi_{a_0}^S(x) = \frac{\bar{f}_{a_0}}{2\sqrt{2N_c}}. \quad (5)$$

The twist-2 LCDA can be expanded in the Gegenbauer polynomials:

$$\Phi_{a_0}(x, \mu) = \frac{1}{2\sqrt{2N_c}} \bar{f}_{a_0}(\mu) 6x(1-x) \sum_{m=1}^{\infty} B_m(\mu) C_m^{3/2}(2x-1), \quad (6)$$

the values for Gegenbauer moments B_1, B_3 have been calculated in [3] as:

$$B_1 = -0.93 \pm 0.10, \quad B_3 = 0.14 \pm 0.08. \quad (7)$$

These values are taken at $\mu = 1 \text{ GeV}$ and the even Gegenbauer moments vanish.

As for the twist-3 distribution amplitudes $\Phi_{a_0}^s$ and $\Phi_{a_0}^T$, they have not been studied in the literature, so we adopt the asymptotic form [8]:

$$\Phi_{a_0}^S = \frac{1}{2\sqrt{2N_c}} \bar{f}_{a_0}, \quad \Phi_{a_0}^T = \frac{1}{2\sqrt{2N_c}} \bar{f}_{a_0} (1-2x). \quad (8)$$

III. THE PERTURBATIVE QCD CALCULATION

Under the two-quark model for the scalar meson a_0 supposition, we would like to use pQCD approach to study B decays into a_0 and π . The decay amplitude can be conceptually written as the convolution,

$$\mathcal{A}(B \rightarrow \pi a_0) \sim \int d^4 k_1 d^4 k_2 d^4 k_3 \text{Tr} [C(t) \Phi_B(k_1) \Phi_\pi(k_2) \Phi_{a_0}(k_3) H(k_1, k_2, k_3, t)], \quad (9)$$

where k_i 's are momenta of light quarks included in each mesons, and Tr denotes the trace over Dirac and color indices. $C(t)$ is the Wilson coefficient which results from the radiative corrections at short distance. In the above convolution, $C(t)$ includes the harder dynamics at larger scale than M_B scale and describes the evolution of local 4-Fermi operators from m_W (the W boson mass) down to $t \sim \mathcal{O}(\sqrt{\bar{\Lambda} M_B})$ scale, where $\bar{\Lambda} \equiv M_B - m_b$. The function $H(k_1, k_2, k_3, t)$ describes the four quark operator and the spectator quark connected by a hard gluon whose q^2 is in the order of $\bar{\Lambda} M_B$, and includes the $\mathcal{O}(\sqrt{\bar{\Lambda} M_B})$ hard dynamics. Therefore, this hard part H can be perturbatively calculated. The function $\Phi_{(\pi, a_0)}$ are the wave functions of π and a_0 , respectively.

Since the b quark is rather heavy we consider the B meson at rest for simplicity. It is convenient to use light-cone coordinate $(p^+, p^-, \mathbf{p}_T)$ to describe the meson's momenta,

$$p^\pm = \frac{1}{\sqrt{2}}(p^0 \pm p^3), \quad \text{and} \quad \mathbf{p}_T = (p^1, p^2). \quad (10)$$

Using these coordinates the B meson and the two final state meson momenta can be written as

$$P_B = \frac{M_B}{\sqrt{2}}(1, 1, \mathbf{0}_T), \quad P_2 = \frac{M_B}{\sqrt{2}}(1, 0, \mathbf{0}_T), \quad P_3 = \frac{M_B}{\sqrt{2}}(0, 1, \mathbf{0}_T), \quad (11)$$

respectively. The meson masses have been neglected. Putting the anti-quark momenta in B , P and S mesons as k_1 , k_2 , and k_3 , respectively, we can choose

$$k_1 = (x_1 P_1^+, 0, \mathbf{k}_{1T}), \quad k_2 = (x_2 P_2^+, 0, \mathbf{k}_{2T}), \quad k_3 = (0, x_3 P_3^-, \mathbf{k}_{3T}). \quad (12)$$

For these considered decay channels, the integration over k_1^- , k_2^- , and k_3^+ in eq.(9) will lead to

$$\mathcal{A}(B \rightarrow \pi a_0) \sim \int dx_1 dx_2 dx_3 b_1 db_1 b_2 db_2 b_3 db_3 \cdot \text{Tr} [C(t) \Phi_B(x_1, b_1) \Phi_\pi(x_2, b_2) \Phi_{a_0}(x_3, b_3) H(x_i, b_i, t) S_t(x_i) e^{-S(t)}], \quad (13)$$

where b_i is the conjugate space coordinate of k_{iT} , and t is the largest energy scale in function $H(x_i, b_i, t)$. In order to smear the end-point singularity on x_i , the jet function $S_t(x)$ [28], which comes from the resummation of the double logarithms $\ln^2 x_i$, is used

$$S_t(x) = \frac{2^{1+2c} \Gamma(3/2 + c)}{\sqrt{\pi} \Gamma(1 + c)} [x(1 - x)]^c, \quad (14)$$

where the parameter $c = 0.4$. The last term $e^{-S(t)}$ in Eq.(13) is the Sudakov form factor which suppresses the soft dynamics effectively [29].

For the considered decays, the related weak effective Hamiltonian $\mathcal{H}_{eff}$ can be written as [30]

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}} \sum_{q=u,c} V_{qb} V_{qd}^* \left[(C_1(\mu) O_1^q(\mu) + C_2(\mu) O_2^q(\mu)) \sum_{i=3}^{10} C_i(\mu) O_i(\mu) \right], \quad (15)$$

with the Fermi constant $G_F = 1.16639 \times 10^{-5} GeV^{-2}$, and the CKM matrix elements V . We specify below the operators in $\mathcal{H}_{eff}$ for $b \rightarrow d$ transition:

$$\begin{aligned} O_1^u &= \bar{d}_\alpha \gamma^\mu L u_\beta \cdot \bar{u}_\beta \gamma_\mu L b_\alpha, & O_2^u &= \bar{d}_\alpha \gamma^\mu L u_\alpha \cdot \bar{u}_\beta \gamma_\mu L b_\beta, \\ O_3 &= \bar{d}_\alpha \gamma^\mu L b_\alpha \cdot \sum_{q'} \bar{q}'_\beta \gamma_\mu L q'_\beta, & O_4 &= \bar{d}_\alpha \gamma^\mu L b_\beta \cdot \sum_{q'} \bar{q}'_\beta \gamma_\mu L q'_\alpha, \\ O_5 &= \bar{d}_\alpha \gamma^\mu L b_\alpha \cdot \sum_{q'} \bar{q}'_\beta \gamma_\mu R q'_\beta, & O_6 &= \bar{d}_\alpha \gamma^\mu L b_\beta \cdot \sum_{q'} \bar{q}'_\beta \gamma_\mu R q'_\alpha, \\ O_7 &= \frac{3}{2} \bar{d}_\alpha \gamma^\mu L b_\alpha \cdot \sum_{q'} e_{q'} \bar{q}'_\beta \gamma_\mu R q'_\beta, & O_8 &= \frac{3}{2} \bar{d}_\alpha \gamma^\mu L b_\beta \cdot \sum_{q'} e_{q'} \bar{q}'_\beta \gamma_\mu R q'_\alpha, \\ O_9 &= \frac{3}{2} \bar{d}_\alpha \gamma^\mu L b_\alpha \cdot \sum_{q'} e_{q'} \bar{q}'_\beta \gamma_\mu L q'_\beta, & O_{10} &= \frac{3}{2} \bar{d}_\alpha \gamma^\mu L b_\beta \cdot \sum_{q'} e_{q'} \bar{q}'_\beta \gamma_\mu L q'_\alpha, \end{aligned} \quad (16)$$

where α and β are the $SU(3)$ color indices; L and R are the left- and right-handed projection operators with $L = (1 - \gamma_5)$, $R = (1 + \gamma_5)$. The sum over q' runs over the quark fields that are active at the scale $\mu = O(m_b)$, i.e., $(q' \in \{u, d, s, c, b\})$.

In the following, we take the $\bar{B}^0 \rightarrow \pi^0 a_0^0$ decay channel as an example to expound. There are 8 type diagrams contributing to this decay, as illustrated in Fig.1. For the factorizable emission diagrams (a) and (b), operators $O_{1-4,9,10}$ are $(V-A)(V-A)$ currents, and the operators O_{5-8} have a structure of $(V-A)(V+A)$, the sum of the their amplitudes are written as $F_{e\pi}$ and $F_{e\pi}^{P1}$. In some other cases, we need to do Fierz transformation for the $(V-A)(V+A)$ operators and get $(S-P)(S+P)$ ones which hold right flavor and color structure for factorization to work. The contribution from the operator $(S-P)(S+P)$ type is written as $F_{e\pi}^{P2}$. Similarly, for the factorizable annihilation diagrams (g) and (h), the contributions from $(V-A)(V-A)$, $(V-A)(V+A)$, $(S-P)(S+P)$ currents are $F_{a\pi}$, $F_{a\pi}^{P1}$ and $F_{a\pi}^{P2}$. For the nonfactorizable spectator diagrams (c, d) and the nonfactorizable annihilation diagrams (e, f), these three kinds of contributions can be written as $M_{e\pi}$, $M_{e\pi}^{P1}$, $M_{e\pi}^{P2}$ and $M_{a\pi}$, $M_{a\pi}^{P1}$, $M_{a\pi}^{P2}$, respectively. Since these amplitudes are similar to those $B \rightarrow f_0(980)K(\pi, \eta^{(\prime)})$ [6, 7] or $B \rightarrow a_0(980)K$ [8], we just need to replace some corresponding wave functions and parameters.

Combining the contributions from different diagrams, the total decay amplitudes for these decays can be written as:

$$\begin{aligned} 2\mathcal{M}(\bar{B}^0 \rightarrow a_0^0 \pi^0) &= \xi_u [(-M_{e\pi} + M_{a\pi} + M_{ea_0} + M_{aa_0})C_2 + (F_{a\pi} + F_{ea_0} + F_{aa_0})a_2] \\ &\quad - \xi_t \left\{ (M_{e\pi}^{P1} + M_{a\pi}^{P1} + M_{ea_0}^{P1} + M_{aa_0}^{P1})(C_5 - \frac{1}{2}C_7) \right. \\ &\quad + (M_{e\pi} + M_{a\pi} + M_{ea_0} + M_{aa_0})(C_3 + 2C_4 - \frac{1}{2}C_9 + \frac{1}{2}C_{10}) \\ &\quad + (M_{e\pi}^{P2} + M_{a\pi}^{P2} + M_{ea_0}^{P2} + M_{aa_0}^{P2})(2C_6 + \frac{1}{2}C_8) + (F_{a\pi} + F_{ea_0} \\ &\quad + F_{aa_0})(2a_3 + a_4 - 2a_5 - \frac{1}{2}a_7 + \frac{1}{2}a_9 - \frac{1}{2}a_{10}) \\ &\quad \left. + (F_{e\pi}^{P2} + F_{a\pi}^{P2} + F_{ea_0}^{P2} + F_{aa_0}^{P2})(a_6 - \frac{1}{2}a_8) \right\}, \end{aligned} \quad (17)$$




FIG. 1: Diagrams contributing to the decay $\bar{B}^0 \rightarrow \pi^0 a_0^0$.

$$\begin{aligned}
\mathcal{M}(\bar{B}^0 \rightarrow a_0^- \pi^+) = & \xi_u [F_{aa_0} a_2 + M_{e\pi} C_1 + M_{aa_0} C_2] - \xi_t \left\{ M_{a\pi}^{P1} \left(C_5 - \frac{1}{2} C_7 \right) \right. \\
& + M_{e\pi}^{P1} (C_5 + C_7) + M_{a\pi} (C_3 + C_4 - \frac{1}{2} C_9 - \frac{1}{2} C_{10}) + M_{aa_0} (C_4 + C_{10}) \\
& M_{e\pi} (C_3 + C_9) + M_{a\pi}^{P2} (C_6 - \frac{1}{2} C_8) + M_{aa_0}^{P2} (C_6 + C_8) \\
& + F_{a\pi} (a_3 + a_4 - a_5 + \frac{1}{2} a_7 - \frac{1}{2} a_9 - \frac{1}{2} a_{10}) \\
& \left. + F_{aa_0} (a_3 + a_9 - a_5 - a_7) + F_{e\pi}^{P2} (a_6 + a_8) + F_{a\pi}^{P2} \left(a_6 - \frac{1}{2} a_8 \right) \right\}, \quad (18)
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}(\bar{B}^0 \rightarrow a_0^+ \pi^-) = & \xi_u [F_{ea_0} a_1 + F_{a\pi} a_2 + M_{ea_0} C_1 + M_{a\pi} C_2] - \xi_t \left\{ M_{aa_0}^{P1} \left(C_5 - \frac{1}{2} C_7 \right) \right. \\
& + M_{ea_0}^{P1} (C_5 + C_7) + M_{aa_0} (C_3 + C_4 - \frac{1}{2} C_9 - \frac{1}{2} C_{10}) + M_{a\pi} (C_4 + C_{10}) \\
& M_{ea_0} (C_3 + C_9) + M_{aa_0}^{P2} (C_6 - \frac{1}{2} C_8) + M_{a\pi}^{P2} (C_6 + C_8) \\
& + F_{aa_0} (a_3 + a_4 - a_5 + \frac{1}{2} a_7 - \frac{1}{2} a_9 - \frac{1}{2} a_{10}) + F_{ea_0} (a_4 + a_{10}) \\
& \left. + F_{a\pi} (a_3 + a_9 - a_5 - a_7) + F_{ea_0}^{P2} (a_6 + a_8) + F_{aa_0}^{P2} \left(a_6 - \frac{1}{2} a_8 \right) \right\}, \quad (19)
\end{aligned}$$

TABLE I: Input parameters used in the numerical calculation[22].

Masses	$m_{a_0} = 0.9847 \text{ GeV},$	$m_0^\pi = 1.3 \text{ GeV},$
	$M_B = 5.28 \text{ GeV},$	$m_\pi = 0.14 \text{ GeV},$
Decay constants	$f_B = 0.19 \text{ GeV},$	$f_\pi = 0.13 \text{ GeV},$
Lifetimes	$\tau_{B^\pm} = 1.671 \times 10^{-12} \text{ s}, \tau_{B^0} = 1.530 \times 10^{-12} \text{ s},$	
CKM	$V_{tb} = 0.9997,$	$V_{td} = 0.0081e^{-i21.6^\circ},$
	$V_{ud} = 0.974,$	$V_{ub} = 0.00393e^{-i60^\circ}.$

$$\begin{aligned}
 \sqrt{2}\mathcal{M}(B^- \rightarrow a_0^0 \pi^-) = & \xi_u [M_{e\pi}C_2 + (-M_{a\pi} + M_{ea_0} + M_{aa_0})C_1 + (-F_{a\pi} + F_{ea_0} + F_{aa_0})a_1] \\
 & - \xi_t \left\{ -M_{e\pi}^{P1}(C_5 - \frac{1}{2}C_7) + (-M_{a\pi}^{P1} + M_{ea_0}^{P1} + M_{aa_0}^{P1})(C_5 + C_7) \right. \\
 & + M_{e\pi}(-C_3 + \frac{1}{2}C_9 + \frac{3}{2}C_{10}) + (-M_{a\pi} + M_{ea_0} + M_{aa_0})(C_3 + C_9) \\
 & + \frac{3}{2}C_8M_{e\pi}^{P2} + (-F_{a\pi} + F_{ea_0} + F_{aa_0})(a_4 + a_{10}) - F_{e\pi}^{P2}(a_6 - \frac{1}{2}a_8) \\
 & \left. + (-F_{a\pi}^{P2} + F_{ea_0}^{P2} + F_{aa_0}^{P2})(a_6 + a_8) \right\}, \quad (20)
 \end{aligned}$$

$$\begin{aligned}
 \sqrt{2}\mathcal{M}(B^- \rightarrow a_0^- \pi^0) = & \xi_u [M_{ea_0}C_2 + (-M_{aa_0} + M_{e\pi} + M_{a\pi})C_1 + F_{ea_0}a_2 + (-F_{aa_0} + F_{e\pi} + F_{a\pi})a_1] \\
 & - \xi_t \left\{ -M_{ea_0}^{P1}(C_5 - \frac{1}{2}C_7) + (-M_{aa_0}^{P1} + M_{e\pi}^{P1} + M_{a\pi}^{P1})(C_5 + C_7) \right. \\
 & + M_{ea_0}(-C_3 + \frac{1}{2}C_9 + \frac{3}{2}C_{10}) + (-M_{aa_0} + M_{e\pi} + M_{a\pi})(C_3 + C_9) \\
 & + \frac{3}{2}C_8M_{ea_0}^{P2} + (-F_{aa_0} + F_{e\pi} + F_{a\pi})(a_4 + a_{10}) - F_{ea_0}^{P2}(a_6 - \frac{1}{2}a_8) \\
 & \left. + F_{ea_0}(-a_4 - \frac{3}{2}a_7 + \frac{3}{2}a_9 + \frac{1}{2}a_{10}) + (-F_{aa_0}^{P2} + F_{ea_0}^{P2} + F_{a\pi}^{P2})(a_6 + a_8) \right\}, \quad (21)
 \end{aligned}$$

where $\xi_u = V_{ub}V_{ud}^*$, $\xi_t = V_{tb}V_{td}^*$. The combinations of the Wilson coefficients are defined as usual [31]:

$$\begin{aligned}
 a_1(\mu) &= C_2(\mu) + \frac{C_1(\mu)}{3}, \quad a_2(\mu) = C_1(\mu) + \frac{C_2(\mu)}{3}, \\
 a_i(\mu) &= C_i(\mu) + \frac{C_{i+1}(\mu)}{3}, \quad i = 3, 5, 7, 9, \\
 a_i(\mu) &= C_i(\mu) + \frac{C_{i-1}(\mu)}{3}, \quad i = 4, 6, 8, 10.
 \end{aligned} \quad (22)$$

IV. NUMERICAL RESULTS AND DISCUSSIONS

In the numerical calculation, we will use the input parameters as listed in Table I.

TABLE II: Branching ratios ($\times 10^{-6}$) for the decays $\bar{B}^0 \rightarrow a_0^0 \pi^0, a_0^\pm \pi^\mp$ and $B^- \rightarrow a_0^- \pi^0, a_0^0 \pi^-$. The first theoretical error is from the the scalar meson decay constant, the second and the third one are Gengebauer moments B_1 and B_3 for twist-2 LCDAs of $a(980)$.

Channel	This work	Data	QCDF [3]
$\bar{B}^0 \rightarrow a_0^0 \pi^0$	$0.51^{+0.08+0.09+0.00}_{-0.07-0.09-0.00}$	—	0.2
$\bar{B}^0 \rightarrow a_0^+ \pi^-$	$0.86^{+0.10+0.14+0.01}_{-0.09-0.14-0.00}$	—	7.6
$\bar{B}^0 \rightarrow a_0^- \pi^+$	$0.51^{+0.05+0.09+0.07}_{-0.06-0.09-0.06}$	—	0.6
$\bar{B}^0 \rightarrow a_0^+ \pi^- + a_0^- \pi^+$	$0.93^{+0.10+0.15+0.02}_{-0.10-0.14-0.00}$	< 3.1	—
$B^- \rightarrow a_0^- \pi^0$	$0.41^{+0.00+0.00+0.00}_{-0.13-0.14-0.12}$	< 1.4	0.2
$B^- \rightarrow a_0^0 \pi^-$	$2.8^{+0.00+0.00+0.00}_{-0.79-0.85-0.58}$	< 5.8	3.4

In the B-rest frame, the decay rates of $B \rightarrow a_0(980)\pi$ can be written as:

$$\Gamma = \frac{G_F^2}{32\pi m_B} |\mathcal{M}|^2 (1 - r_{a_0}^2) \quad (23)$$

where $r_{a_0} = m_{a_0}/m_B$ and $\mathcal{M}$ is the total decay amplitude of $B \rightarrow a_0(980)\pi$, which has been given in section III.

Using the wave functions as specified in previous section and the input parameters listed in Table I, it is straightforward to calculate the CP-averaged branching ratios for the considered decays, which are listed in Table II. In this table, we have included theoretical errors arising from the uncertainties in the scalar meson decay constant $\bar{f}_{a_0}$ and the Gengebauaer moments B_1 and B_3 for twist-2 LCDAs of $a_0(980)$.

From the numerical results, one can find that: Firstly, the branching ratio of $\bar{B}^0 \rightarrow a_0^0 \pi^0$ is larger than that of $\bar{B}^0 \rightarrow f_0(980)\pi^0$ [7], for the small $u\bar{u}$ and $d\bar{d}$ component in the $f_0(980)$, but much smaller than the branching ratio $Br(\bar{B}^0 \rightarrow \pi^0 \pi^0) = (1.62 \pm 0.31) \times 10^{-6}$. Since the scalar meson $a_0(980)$ has vanishing decay constants in the isospin limit, the $a_0^- \pi^+$ rate is smaller than that of its conjugated channel. Secondly, the ratio of these two rates is about 1/2. If one includes the interference between these two decay modes, the branching ratio of $\bar{B}^0 \rightarrow a_0^- \pi^+ + a_0^+ \pi^-$ is close to 1×10^{-6} . Thirdly, for the two charged decays, it is the same reason with the previous two neutral ones, and the $a_0^- \pi^0$ rate is much smaller than that of $a_0^0 \pi^-$. Lastly, the 2-quark model supposition of $a_0(980)$ can not be ruled out by the current experimental data, and this point is different from the QCDF prediction [3]. Since only the upper limits for these channels are available now and the daughter branching fraction has been taken to be 100%, so the further accurate data are needed to clarify this discrepancy.

In Fig. 2, we plot the branching ratios of $\bar{B}^0 \rightarrow a_0^0 \pi^0, a_0^- \pi^+ + a_0^+ \pi^-$ and $B^- \rightarrow a_0^0 \pi^-, a_0^- \pi^0$ as functions of the CKM angle $\alpha = \arg[-\frac{V_{td}^* V_{tb}}{V_{ud}^* V_{ub}}]$. From these figures, it is found that the branching ratio of $\bar{B}^0 \rightarrow a_0^- \pi^+ + a_0^+ \pi^-$ is more sensitive to the angle α than those of other three decay channels. It has been claimed in Ref.[3] that the $B^0 - \bar{B}^0$ interference plays no role in the $a_0^\pm \pi^\mp$ channels. We disagree and we argue that though the $B^0 - \bar{B}^0$ interference in the $a_0^\pm \pi^\mp$ channels may be not as strong as that in some other channels, such as $a_1(1260)^\mp \pi^\pm$ modes. There must exist certain interference among them,




FIG. 2: Branching ratios (in units of 10^{-6}) of (a) $\bar{B}^0 \rightarrow a_0^0 \pi^0$ (solid curve), $\bar{B}^0 \rightarrow a_0^- \pi^+ + a_0^+ \pi^-$ (dashed curve) and (b) $B^- \rightarrow a_0^- \pi^0$ (solid curve), $B^- \rightarrow a_0^0 \pi^-$ (dashed curve) decays as a function of the CKM angle α .

as can be seen easily in the numerical values in Table II, so it may be unreasonable to regard the decays $\bar{B}^0 \rightarrow a_0^\mp \pi^\pm$ as the self-tagging decays in the experiment.

Now we turn to the evaluations of the CP-violating asymmetries of $B^- \rightarrow a_0^- \pi^0$, $a_0^0 \pi^-$ and $\bar{B} \rightarrow a_0^0 \pi^0$, $a_0^\pm \pi^\mp$ decays in pQCD approach. For the charged decay channels, the direct CP-violating asymmetry can be defined as:

$$\mathcal{A}_{CP}^{dir} = \frac{|\bar{\mathcal{A}}|^2 - |\mathcal{A}|^2}{|\mathcal{A}|^2 + |\bar{\mathcal{A}}|^2}. \quad (24)$$

For the neutral decays $\bar{B}^0 \rightarrow a_0^0 \pi^0$, there are both direct CP asymmetry $\mathcal{A}_{CP}^{dir}$ and mixing-induced CP asymmetry $\mathcal{A}_{CP}^{mix}$. The time dependent CP asymmetry of B decay into a CP eigenstate f ($a_0^0 \pi^0$) is defined as:

$$\mathcal{A}_{CP}(t) = \mathcal{A}_{CP}^{dir}(B_d \rightarrow f) \cos(\Delta M t) + \mathcal{A}_{CP}^{mix}(B_d \rightarrow f) \sin(\Delta M t), \quad (25)$$

with

$$\mathcal{A}_{CP}^{dir}(B_d \rightarrow f) = \frac{|\lambda|^2 - 1}{1 + |\lambda|^2}, \quad \mathcal{A}_{CP}^{mix}(B_d \rightarrow f) = \frac{2Im\lambda}{1 + |\lambda|^2}, \quad (26)$$

$$\lambda = \eta e^{-2i\beta} \frac{\mathcal{A}(\bar{B}_d \rightarrow f)}{\mathcal{A}(B_d \rightarrow f)}, \quad (27)$$

where $\eta = \pm 1$ depends on the CP eigenvalue of f , ΔM is the mass difference of the two neutral B meson eigenstates.

As to the decays $\bar{B} \rightarrow a_0^\pm \pi^\mp$, since both B^0 and $\bar{B}^0$ can decay into the final state $a_0^+ \pi^-$ and $a_0^- \pi^+$, the four time-dependent decay widths for $B^0(t) \rightarrow a_0^+ \pi^-$, $\bar{B}^0(t) \rightarrow a_0^- \pi^+$, $B^0(t) \rightarrow a_0^- \pi^+$ and $\bar{B}^0(t) \rightarrow a_0^+ \pi^-$ can be expressed by four basic matrix elements:

$$\begin{aligned} g &= \langle a_0^+ \pi^- | H_{eff} | B^0 \rangle, & h &= \langle a_0^+ \pi^- | H_{eff} | \bar{B}^0 \rangle, \\ \bar{g} &= \langle a_0^- \pi^+ | H_{eff} | \bar{B}^0 \rangle, & \bar{h} &= \langle a_0^- \pi^+ | H_{eff} | B^0 \rangle. \end{aligned} \quad (28)$$




FIG. 3: The direct CP asymmetries (a) of the decays $\bar{B}^0 \rightarrow a_0^0 \pi^0$ (solid curve), $B^- \rightarrow a_0^- \pi^0$ (dashed curve), $B^- \rightarrow a_0^0 \pi^-$ (dotted curve) and the CP asymmetry parameters (b) of the decay $\bar{B}^0 \rightarrow a_0^+ \pi^- + a_0^- \pi^+$: $a_{\epsilon'}$ (dash-dotted curve), $a_{\bar{\epsilon}'}$ (dotted curve), $a_{\epsilon+\epsilon'}$ (dashed curve) and $a_{\epsilon+\bar{\epsilon}'}$ (solid curve) as functions of the CKM angle α .

Following the notation of Ref.[32, 33], the four CP violating parameters are given by the following formulae:

$$\begin{aligned} a_{\epsilon'} &= \frac{|g|^2 - |h|^2}{|g|^2 + |h|^2}, & a_{\epsilon+\epsilon'} &= \frac{-2\text{Im}(\frac{q}{p} \frac{h}{g})}{1 + |h/g|^2} \\ a_{\bar{\epsilon}'} &= \frac{|\bar{h}|^2 - |\bar{g}|^2}{|\bar{h}|^2 + |\bar{g}|^2}, & a_{\epsilon+\bar{\epsilon}'} &= \frac{-2\text{Im}(\frac{q}{p} \frac{\bar{g}}{\bar{h}})}{1 + |\bar{g}/\bar{h}|^2}, \end{aligned} \quad (29)$$

where $q/p = e^{-2i\beta}$ is defined via

$$B_1 = p|B^0\rangle + q|\bar{B}^0\rangle, \quad B_2 = p|B^0\rangle - q|\bar{B}^0\rangle, \quad (30)$$

with $|p|^2 + |q|^2 = 1$ and β being a CKM angle.

From the Fig.3(b), one can find the central values of the CP-violation parameters:

$$a_{\epsilon'} = 0.31, \quad a_{\epsilon+\epsilon'} = 0.94, \quad a_{\bar{\epsilon}'} = 0.93, \quad a_{\epsilon+\bar{\epsilon}'} = 0.32; \quad (31)$$

for $\alpha = 100^\circ$.

The direct CP asymmetries are shown in Fig.3(a). The branching ratio of decay $B^- \rightarrow a_0^0 \pi^-$ is the largest one among the considered channels, while its direct CP asymmetry is the smallest, 14%. Although the CP asymmetries for the decays $\bar{B}^0 \rightarrow a_0^0(980)\pi^0$, $B^- \rightarrow a_0^-(980)\pi^0$ are large, about $(70 \sim 80)\%$, it is still difficult to measure them, since their branching ratios are small, around $(4 \sim 5) \times 10^{-7}$.

V. CONCLUSION

In this paper, we calculate the branching ratios and CP-violating asymmetries of $\bar{B}^0 \rightarrow a_0^0 \pi^0$, $\bar{B}^0 \rightarrow a_0^+ \pi^-$, $\bar{B}^0 \rightarrow a_0^- \pi^+$, $B^- \rightarrow a_0^0 \pi^-$ and $B^- \rightarrow a_0^- \pi^0$ decays in the pQCD

factorization approach by identifying $a_0(980)$ as the 2-quark content. Using the decay constants and light-cone distribution amplitudes derived from QCD sum-rule method, we find that:

- Since the scalar meson $a_0(980)$ has vanishing decay constant in the isospin limit, one can find $Br(\bar{B}^0 \rightarrow a_0^+ \pi^-) > Br(\bar{B}^0 \rightarrow a_0^- \pi^+)$ and $Br(B^- \rightarrow a_0^0 \pi^-) > Br(B^- \rightarrow a_0^- \pi^0)$.
- If one includes the $B^0 - \bar{B}^0$ interference in the $a_0^\pm \pi^\mp$ channels, $Br(\bar{B}^0 \rightarrow a_0^+ \pi^- + a_0^- \pi^+) = 0.93 \times 10^{-6}$, which is close to $Br(\bar{B}^0 \rightarrow a_0^+ \pi^-) = 0.86 \times 10^{-6}$. There must exist certain interference in these two neutral channels.
- Although the CP asymmetries of $\bar{B}^0 \rightarrow a_0^0(980)\pi^0, B^- \rightarrow a_0^-(980)\pi^0$ are large, about $(70 \sim 80)\%$, it is still difficult to measure them, since their branching ratios are small.
- For our predictions agree well with the currently available experimental upper limits, the 2-quark model supposition for $a_0(980)$ can not be ruled out. This point of view is different from the conclusion obtained by using the QCDF approach.

Acknowledgment

Z.Q. Zhang would like to thank H.Y. Cheng and W. Wang for helpful discussions. This work is partly supported by the National Natural Science Foundation of China under Grant No.10575052 and 10735080.

-
- [1] A.K. Giri , B. Mawlong , R. Mohanta Phys. Rev. D **74**, 114001 (2006).
 - [2] H.Y. Cheng ,K.C. Yang, Phys. Rev. D **71**, 054020 (2005).
 - [3] H.Y. Cheng , C.K. Chua , K.C. Yang Phys. Rev. D **73**, 014017 (2006).
 - [4] C.H. Cheng, Phys. Rev. D **67**, 014012 (2003).
 - [5] C.H. Cheng, Phys. Rev. D **67**, 094011 (2003).
 - [6] W. Wang , Y.L. Shen , Y. Li , C.D. Lü Phys. Rev. D **74**, 114010 (2006).
 - [7] Z.Q. Zhang and Z.J. Xiao, Chin. Phys. C**33** (2009) in press; arXiv: 0812.2314 [hep-ph].
 - [8] Y.L. Shen, W. Wang, J. Zhu, C.D. Lu Eur. Phys. J. C **50**:877-887, (2007).
 - [9] D.S. Du, J.W. Li and M.Z. Yang, Phys. Lett. B **619**, 105 (2005).
 - [10] M.Z. Yang, Phys. Rev. D **73**, 034027 (2006).
 - [11] M.Z. Yang, Mod.Phys.Lett.A**21**, 1625 (2006).
 - [12] Belle Collaboration, A. Garmash *et al.*, Phys. Rev. D **65**, 092005 (2002).
 - [13] BaBar Collaboration, B.Aubert *et al.*, Phys. Rev. D **70**, 092001 (2004).
 - [14] Belle Collaboration, A. Garmash *et al.*, Phys. Rev. D **71**, 092003 (2005).
 - [15] J. Dragic , talk presented at the HEP2005 Europhysics Conference in Lisboa, Portugal, July 21-27, 2005.
 - [16] BaBar Collaboration, B. Aubert, *et al.*, Phys. Rev. D **73**, 031101 (2006) ; Phys. Rev. Lett. **94**, 041802 (2005).

- [17] BaBar Collaboration, B. Aubert *et al.*, Phys. Rev. D **72** 072003 (2005).
- [18] BaBar Collaboration, B. Aubert *et al.*, Phys. Rev. D **70** 111102(2004); Phys. Rev. D **75** 111102(2007).
- [19] D. Delepine, J.L. Lucio M., and C.A. Ramirez, Eur. Phys. J. C **45**:693-700(2006).
- [20] BaBar Collaboration, B. Aubert, *et al.*, Submitted to the 33rd International Conference on High-Energy Physics, ICHEP 06, 26 July - 2 August 2006, Moscow, Russia, hep-ex/0607064.
- [21] BaBar Collaboration, B. Aubert, *et al.*, Phys. Rev. D **77**, 011101(2008); Erratum-ibid.**77**, 039903 (2008); Erratum-ibid.**77**, 019904 (2008).
- [22] Particle Data Group, C. Amsler *et al.*, Phys. Lett. B **667**,1(2008).
- [23] F.E. Close, N.A. Törnqvist *et al.*, J. Phys. G **28**, R249 (2002).
- [24] N.A. Tornqvist, Phys. Rev. Lett. **49**, 624 (1982);
- [25] G.L. Jaffe Phys. Rev. D **15**, 267 (1977); Erratum-ibid.Phys. Rev. D **15** 281 (1977); A.L. Kataev, Phys. Atom. Nucl. 68, 567 (2005), Yad. Fiz. 68, 597(2005); A. Vijande, A. Valcarce, F. Fernandez and B. Silvestre-Brac, Phys. Rev. D**72**, 034025 (2005).
- [26] J. Weinstein , N. Isgur , Phys. Rev. Lett. **48**, 659 (1982); Phys. Rev. D **27**, 588 (1983); **41**, 2236 (1990); M.P. Locher et al., Eur. Phys. J. C **4**, 317 (1998).
- [27] V. Baru et al., Phys. Lett. B**586**, 53 (2004).
- [28] H.N. Li, Phys. Rev. D **66**, 094010 (2002).
- [29] H.N. Li and B. Tseng, Phys. Rev. D **57**, 443 (1998).
- [30] G. Buchalla , A.J. Buras , M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
- [31] A. Ali, G. Kramer, C.D. Lü , Phys. Rev. D **58**, 094009 (1998).
- [32] C.D. Lü , M.Z. Yang, Eur. Phys. J. C **23**, 275 (2002).
- [33] Z.Q. Zhang and Z.J. Xiao, Eur. Phys. J. C **59**, 49 (2009).